



Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Medicine

journal homepage: www.elsevier.com/locate/apjtm



Document heading doi:

## First report on natural *Leishmania* infection of *Phlebotomus sergenti* due *Leishmania tropica* by high resolution melting curve method in South–eastern Iran

Aghaei Afshar A<sup>1</sup>, Rassi Y<sup>2\*</sup>, Sharifi I<sup>1\*</sup>, Vatandoost H<sup>2</sup>, Mollaie HR<sup>3</sup>, Oshaghi M.A<sup>2</sup>, Abai MR<sup>2</sup>, Rafizadeh S<sup>4</sup>

<sup>1</sup>Leishmaniasis Research Center, Kerman University of Medical Sciences, Kerman, Iran

<sup>2</sup>Department of Medical Entomology & Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

<sup>3</sup>Department of Medical Virology, Tehran University of Medical Sciences, Tehran, Iran

<sup>4</sup>Department of Deputy of Strategic Planning, Reference Health Laboratories Research Center, Deputy of Treatment, Ministry of Health & Medical Education of Health, Tehran, Iran

### ARTICLE INFO

#### Article history:

Received 9 October 2013

Received in revised form 19 December 2013

Accepted 15 January 2014

Available online 20 February 2014

#### Keywords:

Vector(s)

*Leishmania tropica*

Real-time PCR

Iran

### ABSTRACT

**Objective:** To identify the *Leishmania* species in infected sand flies by Real-time PCR coupled with HRM analysis. **Methods:** Real-time PCR coupled with HRM analysis targeting the first internal transcribed spacer (ITS1) of nuclear ribosomal DNA as the genetic marker was used to identify and distinguish *Leishmania* species in sand flies specimens. **Results:** Three out of 115 females of *Phlebotomus sergenti* (*P. sergenti*) (2.6%) were positive to *Leishmania tropica* (*L. tropica*). **Conclusions:** This is the first report on *P. sergenti* as the main and proven vector of anthroponitic cutaneous leishmaniasis in Dehbakri County using Real-time PCR coupled with HRM analysis. This method is rapid, sensitive and specific for diagnosing of parasites in infected Sand flies and ideal for large scale genotyping projects.

## 1. Introduction

Cutaneous leishmaniasis (CL) is endemic in many parts of the world and considered as a major public health problem [1]. The disease is one of the main health problem in Iran. Zoonotic cutaneous leishmaniasis (ZCL) due to *Leishmania major* (*L. major*) is endemic in 50% of the 31 provinces of Iran, whereas anthroponitic cutaneous leishmaniasis (ACL) associated with *Leishmania tropica* (*L. tropica*) is an old endemic disease in 8 provinces, especially in South East Iran [2–6].

Two epidemiological forms of CL occur in Bam district, ACL is mainly limited to the city, with its main reservoir of host of human. Anthroponitic Cutaneous Leishmaniasis is a serious problem due to its difficulty of treatment and longer duration, hence potentially severe long-term complications [7–9]. However ZCL is the more prevalent form. After the massive earthquake of Bam in December 2003 the prevalence of CL cases increased among school children. It increased from 2% in 2005 to 5% in 2008 (unpublished data). The disease is outspread and a new emerging focus of cutaneous leishmaniasis was confirmed in Dehbakri County, adjacent to Bam. According to published data, this paper is the first report on identifying of responsible sand flies to transmitting of disease agent to human, employing Real-time PCR coupled with HRM analysis.

In many areas, however, despite considerable research on these diseases, the main ‘reservoir’ hosts and the species of sand fly responsible for most transmission have still to be

\*Corresponding author: Professor Yavar Rassi, Department of Medical Entomology & Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

E-mail: rassy@tums.ac.ir

Professor Iraj Sharifi, Leishmaniasis Research Center, Kerman University of Medical Sciences, Kerman, Iran.

E-mail: iraj.sharifi@yahoo.com

identified. In many foci of CL there is at least one species of sand fly that is sufficiently common and anthropophilic to be considered a probable vector, although good evidence to support this belief, such as the detection, in wild caught females of this species, of the parasites causing the CL, is lacking. The prevalence of infection in even the primary vector may be quite low, making the detection of sand fly infection, and particularly the detection of any temporal trends in the prevalence of sandfly infection, difficult, especially if dissection and microscopy constitute the detection method. Furthermore this traditional methods do not discriminate the organism in species level and often with variable and low sensitivity<sup>[10,11]</sup>.

Molecular methods are currently being employed to detect *Leishmania* infection up to genus, complex or species level<sup>[12,13]</sup>. High-resolution melting (HRM) analysis is a relatively new technique that allows direct characterization of PCR amplicons in a closed system. It measures changes in the fluorescence intensity of a DNA intercalating dye during dissociation from double-stranded DNA to single-stranded DNA and can differentiate between single nucleotide polymorphisms (SNP)<sup>[14]</sup>.

Real-time PCR offers several advantages over traditional PCR, including faster processing time, higher sensitivity and decreased contamination risk. A previous study used HRM PCR assay for Old World *Leishmania* based on gene 7SL-HRM that can be used to differentiate between *L. tropica*, *L. major*, and the *Leishmania donovani* complex<sup>[15]</sup>.

PCR processing. Nowadays, the HRM has mostly been used in human clinical studies<sup>[16–20]</sup>. However, the application of the HRM technique to the diagnosis of parasitic organisms has been rather limited and the method has mostly been applied in molecular studies of parasitic protozoa such as the old world *Leishmania* spp<sup>[14]</sup>. The aim of the present study was to evaluate the potential use of ITS1 gene for identifying Old World *Leishmania* species by HRM method based real-time PCR. The method was successfully applied to identify *Leishmania* species in sand flies collected from Dehbakri county, Bam, South eastern Iran.

## 2. Material and methods

### 2.1. Study area

This study was carried out in Dehbakri county, Bam district, Kerman province during summer of 2011. Dehbakri is located 60 km west of Bam city. This area has a semi-desert climate temperate in summer and cold at winter. The mean of monthly maximum and minimum temperatures were 40 °C and –5 °C in July and Dec, respectively. The total annual rainfall was 220 mm. The minimum and maximum of monthly relative humidity were 45% and 92%, respectively in

July and January.

### 2.2. Sample collection and identification

Sand flies were caught using aspirator from 8.00 pm till 2.00 am during summer 2011 where cases of ACL had been reported. The caught sand flies were transferred to the entomological cage and then were kept by wet towel and transported to laboratory research center for identification and Detection of *Leishmania* species. All the collected sand flies were mounted separately using Puri's media for species identification. Male and female were considered separately<sup>[21–23]</sup>.

### 2.3. DNA extraction

Sand flies were washed in 1% detergent (washing-up liquid) solution for 5 min, Heads and last abdominal segments were kept for morphological identification based on the keys described by Leger *et al.*, 1986<sup>[24]</sup>. Individual code names, consisting of a letter(s) taken from the collection area name followed by a number were assigned. DNA was extracted by maceration in a micro tube using a plastic pestle, followed by addition of 35  $\mu$  L of lysis buffer (100 mM TRIS-HCl, 100 mM NaCl, 25 mM EDTA, 0.5% SDS, pH 8.0) and another maceration step. The test samples were digested overnight at 37 °C by proteinase K (1.25  $\mu$  L of a 10 mg/mL solution) and the DNA extracted by phenol-chloroform. The DNA pellet was resuspended in 20  $\mu$  L of TE (10 mM TRIS-HCl pH 8.0, 1 mM EDTA). Ten  $\mu$  L were used to estimate the DNA concentration and purity at 280 and 260 nm in a spectrophotometer, then discarded. The remaining 10  $\mu$  L were stored at –20 °C until use. Double distilled water and DNA from *L. major* and *L. tropica* provided to the Iran Institute of Pasteur by the World Health Organization were used as negative and positive controls.

### 2.4. HRM-real-time PCR assay

We designed two primers AGCTGGATCATTTCGGATG and ATCGCGACACGTTATGTGAG using the software Beacon as forward and reverse respectively. Real-time PCR was performed in a total reaction mixture of 20  $\mu$  L containing 10  $\mu$  L of HRM Master Mix, 10 pmole of each primer, approximately 10 ng/mL of genomic DNA and sterile deionized water using a 7500 fast real-time PCR system. Infection was detected by internal transcribed spacer 1 (ITS1) real-time PCR and high-resolution melt analysis (ITS1-HRM) PCR. All samples were tested in duplicates and results were compared with those from HRM analysis of positive controls for each assay. Positive samples also were verified by kDNA PCR.


### 3. Results

#### 3.1. Sand fly identification


A total of 1 261 sand flies were captured and identified to the species level. Fourteen species including, 5 species of *Phlebotomus* genus and 9 species of *Sergentomyia* genus were identified. They were *Phlebotomus sergenti* (*P. sergenti*), *Phlebotomus papatasi* (*P. papatasi*), *Phlebotomus monglensis*, *Phlebotomus major*, *Phlebotomus longiductus*, *Sergentomyia sintoni*, *Sergentomyia baghdadis*, *Sergentomyia sumbarica*, *Sergentomyia africana*, *Sergentomyia dentata*, *Sergentomyia squemipleuris*, *Sergentomyia grekovi*, *Sergentomyia mervynae* and *Sergentomyia antennata*. Two species of *P. sergenti* (67.8%) and *P. papatasi* (19.4) were the dominant specimens respectively.

#### 3.2. *Leishmania* species identification

We tested 164 specimens of sand flies including 115 *Ph. sergenti* and 49 *P. papatasi* by ITS1–HRM PCR. The overall *Leishmania* infection rate for *P. sergenti* was 2.6% (3/115) and all *P. papatasi* samples were negative. Positive specimens by ITS1–HRM PCR were also positive by kDNA PCR and produced a 750 bp kDNA product. Infected sand flies had been collected from indoors places with empty position of abdomen. This is the first report on natural infection of *P. sergenti* to *L. tropica* using high resolution melting curve (HRM) method in Iran. According our results this species was the only infected sand fly and it seems was responsible to transmitting of *L. tropica* among human.



**Figure 1.** Differentiation *L. tropica* positive control and *L. major* positive control using HRM method.



**Figure 2.** Representative curve of the melting curves of ITS–1 amplicons for infected *P. sergenti* species to *L. tropica*, Dehbakri County, Southeastern Iran, 2011.

### 4. Discussion

Control of leishmaniasis need to understand on ecology and epidemiology of disease in endemic areas. There is a major problem for epidemiologists both in identification of reservoir hosts and in detection of vectors. Therefore, finding naturally infected sand flies is essential in identifying a species as a vector of *Leishmania* and in studying infection rates in areas of endemicity[25, 26].

The applicability of kDNA for the detection and identification of *Leishmania* within sand flies by DNA hybridization have been shown previously[27]. In recent years, several molecular techniques, mainly those based on PCR have been developed for the specific identification and characterization of *Leishmania* species[28–30]. Although other techniques are sensitive and specific for identification of *Leishmania* species, they are laborious and time consuming, especially the post–PCR processing steps. In addition, there is also a higher risk of contamination, more expensive (*e.g.*, DNA sequencing) and the techniques only provide qualitative information. A highly sensitive method is needed to determine sand fly infection of *Leishmania* parasite. The choice of the gene used for HRM analysis is important in developing a successful assay, since it can take advantage of small differences in melting curves to distinguish between organisms with highly homologous sequences[14].

The ITS1 gene can be used for diagnosis of human leishmaniasis, as well as epidemiological studies on potential reservoir hosts, sand fly vectors, and parasite genotypes[12]. In the present study, we have successfully utilized HRM analysis along with real–time PCR for the rapid detection and species identification of *Leishmania* species in sand flies samples. To the best of our knowledge, this is the first report on the utilization of the HRM approach for rapid detection and discrimination of sand flies species by employing the ITS1 of nuclear ribosomal DNA as a genetic marker in Iran. We report *L. tropica* infections in *P. sergenti*(2.6%) specimens, the only proven vector of *L. tropica* in Iran.

In conclusion, using HRM assay for identifying of Old World *Leishmania* species is rapid, sensitive, specific, and simple and can be used to directly diagnose parasites in the sand flies vector with a minimum of operator post–PCR manipulation and interpretation.

#### Conflict of interest statement

We declare that we have no conflict of interest.

#### Acknowledgments

This investigation was supported by the Iranian Center of Diseases Management, Ministry of Health and Medical

Education. We wish to express our sincere thanks for providing facilities needed for this study. The authors would like to thank the Leishmaniasis Research Center, Kerman University of Medical Sciences and the School of Public Health, Tehran University of Medical Sciences for their close collaboration. This study was financially supported by School of Public Health, Tehran University of Medical Sciences, and Leishmaniasis Research Center, Kerman University of Medical Sciences, project No.10487

## References

- [1] Desjeux P. Leishmaniasis: current situation and new perspectives. *Comp Immunol Microbiol Infect Dis* 2004; **27**: 305–318.
- [2] Rassi Y, Javadian E, Jalali M, Motazedian MH, Vatandoost H. Investigation on zoonotic cutaneous leishmaniasis, southern Iran. *Iran J Public Health* 2004; **33**(1): 31–35.
- [3] Rassi Y, Javadian E, Amin M, Rafizadeh S, Vatandoost H, Motazedian H. *Meriones libycus* is the main reservoir of zoonotic cutaneous leishmaniasis in south Islamic Republic of Iran. *East Med Health* 2006; **12**(3–4): 475–457.
- [4] Rassi Y, Gassemi MM, Javadian E, Rafizadeh S, Motazedian H, Vatandoost H. Vectors and reservoirs of cutaneous leishmaniasis in Marvdasht district, southern Islamic Republic of Iran. *Est Mediterr Health J* 2007; **3**(3): 686–693.
- [5] Rassi Y, Oshaghi MA, Mohammadi Azani S, Abaie MR, Rafizadeh S, et al. Molecular detection of *Leishmania* infection due to *Leishmania major* and *Leishmania turanica* in the vectors and reservoir host in Iran. *Vector-borne Zoonotic Dis* 2011; **11**: 145–150.
- [6] Nadim A. *Epidemiology of leishmaniasis in Iran, leishmania parasites and leishmaniasis*. Tehran, Iran: University press; 2008, p. 191–211.[In Persian]
- [7] Aghaei Afshar A, Rassi Y, Sharifi I, Abai MR, Vatandoost H, Oshaghi MA, et al. Susceptibility status of *Phlebotomus papatasi* and *P. sergenti* (Diptera: Psychodidae) to DDT and Deltamethrin in a focus of cutaneous leishmaniasis after earthquake strike in Bam, Iran. *J Arthropod-Borne Dis* 2011; **5**(2): 32–41.
- [8] Sharifi I, Zemani F, Aflatoonian MR, Fekri AR. An epidemic of cutaneous leishmaniasis in Baft district in Kerman Province and its probable causative risk factors. *Iranian J Epidemiol* 2008; **4**(1): 53–58.
- [9] Sharifi I, Poursmaelian S, Aflatoonian MR, Ardakani RF, Mirzaei M, Fekri AR, et al. Emergence of a new focus of anthroponotic cutaneous leishmaniasis due to *Leishmania tropica* in rural communities of Bam district after the earthquake, Iran. *Trop Med Int Health* 2011; **16**(4): 510–513.
- [10] Hajjarian H, Vasigheh F, Mohebbali M, Rezaei S, Mamishi S, Charedar S. Direct diagnosis of *Leishmania* species on serosity materials punctured from cutaneous leishmaniasis patients using PCR-RFLP. *J Clin Lab Anal* 2011; **25**(1): 20–24.
- [11] Shahbazi F, Shahabi S, Kazemi B, Mohebbali M, Abadi AR, Zare Z. Evaluation of PCR assay in diagnosis and identification of cutaneous leishmaniasis: a comparison with the parasitological methods. *Parasitol Res* 2008; **103**(5): 1159–1162.
- [12] Bensoussan-Hermano E, Nasereddin A, Jonas F, Schnur LF, Jaffe CL. Comparison of PCR assays for diagnosis of cutaneous leishmaniasis. *J Clin Microbiol* 2006; **44**: 1435–1439.
- [13] Nasereddin A, Bensoussan-Hermano E, Schoniane G, Baneth G, Jaffe CL. Molecular diagnosis of Old World cutaneous leishmaniasis and species identification by use of a reverse line blot hybridization assay. *J Clin Microbiol* 2008; **46**: 2848–2855.
- [14] Talmi-Frank D, Nasereddin A, Schnur LF, Schonian G, Töz SO. Detection and identification of Old World *Leishmania* by high resolution melt analysis. *PLoS Negl Trop Dis* 2010; **4**(1): e581.
- [15] Nasereddin A, Jaffe CL. Rapid diagnosis of Old World *Leishmaniasis* by high-resolution melting analysis of the 7SL RNA gene. *J Clin Microbiol* 2010; **2240**–2242.
- [16] Wittwer CT, Reed GH, Gundry CN, Vandersteen JG, Pryor RJ. High resolution genotyping by amplicon melting analysis using LC Green. *Clin Chem* 2003; **49**: 853–860.
- [17] Zhou L, Vandersteen J, Wang L, Fuller T, Taylor M. High-resolution DNA melting curve analysis to establish HLA genotypic identity. *Tissue Antigens* 2004; **64**: 156–164.
- [18] Liew M, Pryor R, Palais R, Meadows C, Erali M. Genotyping of single-nucleotide polymorphisms by high-resolution melting of small amplicons. *Clin Chem* 2004; **50**: 1156–1164.
- [19] Radvansky J, Resko P, Surovy M, Minarik G, Ficek A. High-resolution melting analysis for genotyping of the myotonic dystrophy type I associated Alu insertion/deletion polymorphism. *Anal Biochem* 2010; **398**: 126–128.
- [20] Saitsu H, Kato M, Okada I, Orii KE, Higuchi T. STXBPI mutations in early infantile epileptic encephalopathy with suppression-burst pattern. *Epilepsia* 2010; **51**: 2397–2405.
- [21] Rassi Y, Hanafi-Bojd AA. Phlebotomine sand flies, vectors of Leishmaniasis: Morphology, biology, ecology, and field and laboratory methods with pictorial key of Iranian sand flies. Noavaran-Elm Publication 2006. (In Persian)
- [22] Lewis DJ. A taxonomic review of the genus *Phlebotomus*. *Bull Br Mus* 1982; **45**: 121–209.
- [23] Theodor O, Mesghali A. On the phlebotominae of Iran. *J Med Entomol* 1964; **1**: 285–300.
- [24] Leger NB, Pesson G, Madulo-Leblond G. Les phlebotomes de Grece. *Biol Gallo-Hellenica* 1986; **11**: 165–192.
- [25] Killick-Kendrick R. Phlebotomine vectors of the leishmaniasis: a review. *J Med Vet Entomol* 1990; **4**: 1–24.
- [26] Sharma U, Singh S. Insect vectors of *Leishmania*: distribution, physiology and their control. *J Vector Borne Dis* 2008; **45**: 255–272.
- [27] Ready PD, Lainson R, Shaw JJ, Souza AA. DNA probes for distinguishing *Psychodopygus wellcomi* from *Psychodopygus complexus* (Diptera: Psychodidae). *Memorias do Instituto Oswaldo Cruz* 1991; **86**: 41–49.
- [28] Nicolas L, Milon G, Prina E. Rapid differentiation of Old World *Leishmania* species by LightCycler polymerase chain reaction and melting curve analysis. *J Microbiol Methods* 2002; **51**(3): 295–299.
- [29] Nicolas L, Prina E, Lang T, Milon G. Real-time PCR for detection and quantitation of *Leishmania* in mouse tissues. *J Clin Microbiol* 2002; **40**(5): 1666–1669.
- [30] Rotureau B, Joubert M, Clyti E, Djossou F, Carme B. *Leishmaniasis* among gold miners, French Guiana. *Emerg Infect Dis* 2006; **12**(7): 1169–1170.